



Last Updated Mar.2012

Bighead knapweed (*Centaurea macrocephala*)

aka Armenian basketflower, globe centaurea, golden thistle, yellow/lemon fluff knapweed, yellow hardhead

Provincial Designation: Prohibited Noxious

Overview:

Bighead knapweed is a long-lived perennial herb native to the Caucasus region of Eastern Europe. It has been a cultivated ornamental for over 200 years. Introduced to the UK and North America in the early 1800's, it is still widely available in the horticultural trade as seed, plants, and cut/dried flowers. This knapweed reproduces by seed but may be propagated by diving the crown. Its tap root develops a woody crown with age. This is the largest *Centaurea* in cultivation or naturalized, so is easily identified.¹

Bighead knapweed flowers July to September.³ Mature seed heads open to disperse the seed. Although seeds are plumed, they are too heavy to be wind dispersed but can attach to animal hair/fur.¹

Bighead knapweed needs disturbance to establish but is then difficult to control. Eradication efforts



PHOTO: Dave Hayman

of a 4 ha infestation in Washington state have been ongoing for about 25 years.¹ It competes with native vegetation for nutrients and light.

Habitat:

It requires loamy, well drained soils with neutral pH and full sun. It is drought tolerant and prefers warm temperate or continental climates with dry summers.¹

Identification:

Stems: Stems are several to many erect, unbranched or sparingly branched near the tips, growing 50-170 cm tall. They are covered with long, soft hairs, which are slightly cobwebby.¹

Leaves: Leaves are lance shaped, 10-30 cm long, and slightly undulating with pointed tips. Basal and lower stem leaves have short petioles (leaf stem). Leaf margins are smooth or shallowly dentate. Upper stem leaves are 5-10 cm long, sessile (no petioles), and slightly undulate with smooth margins. Leaves are covered with short hairs and are dotted with resin glands.²

Flowers: Flowers are disc-shaped, borne singly and composed of many yellow florets, with clusters of reduced leaves directly below the flower heads.¹ Involucre (flower base) are 25-35 mm diameter and covered with 1-2 mm wide brown bracts having fringed edges, sometimes with weak spines. Seeds are indehiscent, 7-8 mm long and with flattened bristles 5-8 mm long.²

Prevention:

Many infestations result from garden escapes – do not grow as an ornamental. Bighead knapweed used in dried flower arrangements facilitates seed dispersal.¹

Control:

Grazing: Cattle will consume bighead



PHOTO: Richard Old, XID Services, Inc., Bugwood.org

knapweed but this is not considered a control method.¹ *Invasive plants should never be considered as forage.*

Mechanical: Repeated mowing will suppress seed production but plants will re-grow.³ Repeated cultivation, where suitable, is an effective control method. Small patches can be hand pulled or dug out but the entire taproot must be removed to prevent re-sprouting.¹

Chemical: Currently there are no herbicides registered for use on bighead knapweed. Always check product labels to ensure the herbicide is registered for use on the target plant in Canada by the Pesticide Management Regulatory Agency. Consult your local Agricultural Fieldman or Certified Pesticide Dispenser for more information.

Biological: None researched to date.

REFERENCES

- ¹ *Centaurea macrocephala*. Invasive Species Compendium. CABl. www.cabi.org
- ² *Centaurea macrocephala* in Flora of North America. www.efloras.org
- ³ Bighead Knapweed. King County Noxious weed control Program Weed Alert. www.kingcounty.gov/weeds